


## IgG vs IgE: What is the Difference?

There is a great deal of misunderstanding about the differences between a sensitivity versus an allergic reaction to food. This bulletin has been created in order to help demystify the confusion between the different types of reactions and tests available.

There are essentially 3 different types of reactions to food in which the immune system inappropriately identifies the food as foreign and mounts an immune response.

1. The autoimmune response – An autoimmune response can give rise to multifactorial symptoms and, if undiagnosed, results in permanent damage. The most commonly known condition in this category is coeliac disease, where there is an autoimmune response to gluten and gliadin. Testing for this disease is typically done through blood work and/or small bowel biopsy. Here laboratory tests focus on the measurement of IgA and IgG class antibodies to the enzyme Tissue Transglutaminase (tTG).
2. A Type I immediate IgE hypersensitivity immune response – This response is more commonly known as a food allergy. This reaction typically occurs very quickly (minutes to hours) after eating an offending food and symptoms range from mild to severe. Many are aware of food allergies causing anaphylaxis (the most common example being peanuts) but they may also experience itchy lips/tongue/throat, stuffy nose, headaches, stomach pain, diarrhoea, gas, bloating, skin reactions, and sudden fatigue. Most people often know their 'trigger foods' because when the reaction occurs it is so quick and occurs soon after eating food. However, sometimes it isn't always obvious, particularly with people who have multiple allergies. In these instances, testing can be very useful and is done by measuring specific IgE antibodies in the blood to the suspected foods and food panels or by skin prick test.
3. Non-immediate (delayed) IgG response – This type of response is referred to as a food sensitivity. This differs from an immediate IgE food allergy response in that symptoms may not show up for hours or days. Due to the delayed response, determining exactly which food(s) are causing symptoms in the body is difficult, especially as people tend to eat the same food(s) every day. Symptoms of food sensitivities include headaches, stomach pain, constipation, diarrhoea, bloating and many more.

*NB: Having a food allergy doesn't mean that a person has a food sensitivity and having a food sensitivity doesn't mean that the patient has an allergy*

Testing for IgG food sensitivities can be confusing for many healthcare providers as this type of reaction is neither obvious nor life-threatening. However, food sensitivities are gaining public and academic awareness as increasingly practitioners and clients are removing offending food groups in order to help support various conditions such as migraines and IBS, helping to reduce inflammation in the body with great success. Unfortunately though, there is some conflicting evidence on the comparison of IgG versus IgE and the testing behind it. Keep in mind that they are different ‘branches’ of the immune system and being allergic to a food doesn’t mean the patient is sensitive to that food and vice versa.

Some negative press and studies conclude that IgG sensitivity testing has no validity or value. However, when we and others have reviewed all the research against food sensitivity testing, the studies routinely point out that when someone withdraws a particular food based on IgG testing results symptoms can improve.

The food sensitivity test therefore serves as a valuable guide to help patients determine which foods to rotate or eliminate out of the diet before re-introducing them at a later date.

Whilst there is no gold standard for IgG testing, in clinical experience food sensitivity testing via the measurement of IgG antibodies can be an effective, simple to perform and well tolerated tool. Providers have been performing such tests worldwide for many years with positive outcomes. Results help guide patients in a reasonable way that makes sense, to remove certain food(s) in order to help resolve a contributing cause of some of their often chronic symptoms.



## Characteristics of IgE- & IgG- Mediated Reactions to Food

| IgE-Mediated - Allergy                                    | IgG-Mediated - Sensitivity                                                                         |
|-----------------------------------------------------------|----------------------------------------------------------------------------------------------------|
| Incidence is relatively low                               | Incidence is relatively high                                                                       |
| Result from infrequent exposure                           | Result from frequent exposure                                                                      |
| Common in children, rare in adults                        | Most common form of food sensitivity reaction in both children & adults                            |
| Very predictable short-term symptoms                      | Chronic, variable symptoms                                                                         |
| Offending food is usually obvious                         | Offending food frequently hard to identify                                                         |
| Basophil/Mast Cell triggered reactions                    | Immune complex trigger                                                                             |
| Histamine/Leukotriene release                             | Inflammatory response                                                                              |
| Patient aware of offending food                           | Patient rarely aware of offending food                                                             |
| Antibody persistent for years                             | Antibody declines over months                                                                      |
| In-vitro or skin prick testing for serum IgE confirmation | In-vitro testing for serum IgG shows reactive food and degree of gut permeability support required |
| Permanent food avoidance and immunotherapy                | Eliminate then rotate food(s), heal gut, improve digestion                                         |

## Allergy vs Sensitivity

| Category         | Allergy                       | Sensitivity                                        |
|------------------|-------------------------------|----------------------------------------------------|
| Food             | e.g. peanut, shellfish        | e.g. gluten, dairy                                 |
| Mediated         | IgE antibodies                | IgG antibodies                                     |
| Rate of response | Immediate after ingestion     | Delayed up to 72 hrs after ingestion               |
| Mechanism        | Rapid production of histamine | Gradual formation of Ag/Ab complexes               |
| Symptoms         | Classical 'allergic' response | Many symptoms affecting any part of the body       |
| Severity         | Can be fatal                  | Not life-threatening                               |
| Permanence       | Can last lifetime             | Can be reversed or reduced by elimination of foods |
| Diagnosis        | Often self-diagnosed          | Rarely self-diagnosed                              |
| Skin-prick test  | Positive                      | Negative                                           |